

Abelian complex structures and related geometries

María Laura Barberis



Universidad Nacional de Córdoba, Argentina
CIEM - CONICET



Encuentro de Geometría Diferencial
Rosario 2012

Joint work with A. Andrada and I. Dotti

- Abelian complex structures.
- Relation to HKT geometry.
- Affine Lie algebras and abelian double products.
- Kähler Lie algebras with abelian complex structures.
- The first canonical connection.

Abelian complex structures

A complex structure J on a real Lie algebra \mathfrak{g} is called *abelian* when it satisfies:

$$[Jx, Jy] = [x, y], \quad \forall x, y \in \mathfrak{g}. \quad (1)$$

Equivalently, $\mathfrak{g}^{1,0}$ is an abelian subalgebra of $\mathfrak{g}^{\mathbb{C}}$.

If G is a Lie group with Lie algebra \mathfrak{g} these conditions imply the vanishing of the Nijenhuis tensor on the invariant almost complex manifold (G, J) , that is, J is integrable on G .

- A *hyperhermitian* structure on a smooth manifold M is $(\{J_\alpha\}_{\alpha=1,2,3}, g)$, where
 - ① $\{J_\alpha\}_{\alpha=1,2,3}$ are complex structures such that
$$J_1 J_2 = -J_2 J_1 = J_3,$$
 - ② g is a Riemannian metric which is Hermitian with respect to J_α , $\alpha = 1, 2, 3$.
- Given a hyperhermitian structure $(\{J_\alpha\}_{\alpha=1,2,3}, g)$ on M , g is called *hyper-Kähler with torsion* (HKT) if there exists a connection ∇ on M satisfying
 - ① $\nabla g = 0$, $\nabla J_\alpha = 0$, $\alpha = 1, 2, 3$,
 - ② the torsion tensor $c(X, Y, Z) = g(X, T(Y, Z))$ is skew-symmetric.

This class of metrics has been introduced by P.S. Howe - G.Papadopoulos (1996).

- Dotti - Fino (2002): If G is a 2-step nilpotent Lie group with a left invariant HKT structure $(\{J_\alpha\}_{\alpha=1,2,3}, g)$, then the hypercomplex structure is abelian.
- B - Dotti - Verbitsky (2009): Let $(N, \{J_\alpha\}_{\alpha=1,2,3}, g)$ be an HKT nilmanifold such that $\{J_\alpha\}$ is left invariant. Then the hypercomplex structure $\{J_\alpha\}$ is abelian.

Properties of Lie algebras carrying abelian complex structures

Let J be an abelian complex structure on the Lie algebra \mathfrak{g} . Then:

- (i) The center \mathfrak{z} of \mathfrak{g} is J -stable.
- (ii) For any $x \in \mathfrak{g}$, $\text{ad}_{Jx} = -\text{ad}_x J$.
- (iii) $\mathfrak{g}' = [\mathfrak{g}, \mathfrak{g}]$ is abelian, equivalently, \mathfrak{g} is 2-step solvable [Petravchuk, 1988].
- (iv) $J\mathfrak{g}'$ is an abelian subalgebra.
- (v) $\mathfrak{g}' \cap J\mathfrak{g}' \subseteq \mathfrak{z}(\mathfrak{g}'_J)$, where $\mathfrak{g}'_J := \mathfrak{g}' + J\mathfrak{g}'$.

- Let (\mathcal{A}, \cdot) be a finite dimensional associative, commutative algebra. Set $\mathfrak{aff}(\mathcal{A}) := \mathcal{A} \oplus \mathcal{A}$ with Lie bracket:

$$[(a, a'), (b, b')] = (0, a \cdot b' - b \cdot a'), \quad a, b, a', b' \in \mathcal{A},$$

In particular, when $\mathcal{A} = \mathbb{R}$ or $\mathcal{A} = \mathbb{C}$, we obtain the Lie algebra of the group of affine motions of either \mathbb{R} or \mathbb{C} .

- Let J be the endomorphism of $\mathfrak{aff}(\mathcal{A})$ defined by

$$J(a, a') = (a', -a), \quad a, a' \in \mathcal{A}.$$

J defines an abelian complex structure on $\mathfrak{aff}(\mathcal{A})$, which we will call standard.

Abelian double products

[Andrada-Salamon, 2005] Consider a finite dimensional real vector space \mathcal{A} with two structures of commutative associative algebra, (\mathcal{A}, \cdot) and $(\mathcal{A}, *)$, with the following compatibility conditions:

$$a * (b \cdot c) = b * (a \cdot c), \quad a \cdot (b * c) = b \cdot (a * c), \quad (2)$$

for every $a, b, c \in \mathcal{A}$.

Then, $\mathcal{A} \oplus \mathcal{A}$ with the bracket:

$$[(a, a'), (b, b')] = (-(a * b' - b * a'), a \cdot b' - b \cdot a'), \quad a, b, a', b' \in \mathcal{A},$$

is a Lie algebra denoted by $(\mathcal{A}, \cdot) \bowtie (\mathcal{A}, *)$ and the endomorphism J defined by

$$J(a, a') = (-a', a), \quad a, a' \in \mathcal{A}, \quad (3)$$

is an abelian complex structure, called the *standard* complex structure on $(\mathcal{A}, \cdot) \bowtie (\mathcal{A}, *)$.

More examples

We show next that there is a large family of Lie algebras with abelian complex structure which are not abelian double products.

Let $\mathfrak{g} = \mathfrak{a} \oplus \mathfrak{v}$ where $\mathfrak{a} = \text{span}\{f_1, f_2\}$ and \mathfrak{v} is a $2n$ -dimensional real vector space. We fix an endomorphism J of \mathfrak{g} such that $J^2 = -I$, $Jf_1 = f_2$ and \mathfrak{v} is J -stable. Given a linear isomorphism T of \mathfrak{v} commuting with $J|_{\mathfrak{v}}$, we define a Lie bracket on \mathfrak{g} such that \mathfrak{a} is an abelian subalgebra, \mathfrak{v} is an abelian ideal and the bracket between elements in \mathfrak{a} and \mathfrak{v} is given by:

$$[f_1, v] = TJ(v), \quad [f_2, v] = T(v), \quad \text{for every } v \in \mathfrak{v}.$$

It turns out that J is an abelian complex structure on \mathfrak{g} .

The Lie algebra \mathfrak{g} is not an abelian double product, unless $n = 1$. In this case, $\mathfrak{g} = \mathfrak{aff}(\mathbb{C})$ with an abelian complex structure which is NOT the standard one.

Theorem (Andrada -B - Dotti, 2011)

Let \mathfrak{g} be a solvable Lie algebra with an abelian complex structure J such that \mathfrak{g} admits a vector space decomposition $\mathfrak{g} = \mathfrak{u} + J\mathfrak{u}$.

Then:

- (i) if \mathfrak{u} is an abelian subalgebra of \mathfrak{g} then $\mathfrak{g} = \mathfrak{a} \oplus J\mathfrak{a}$ is an abelian double product with $\mathfrak{a} \subset \mathfrak{u}$;
- (ii) if \mathfrak{u} is an abelian ideal of \mathfrak{g} and, moreover, $\mathfrak{g}' \cap J\mathfrak{g}' = \{0\}$, then (\mathfrak{g}, J) is holomorphically isomorphic to $\text{aff}(\mathcal{A})$ for some commutative associative algebra (\mathcal{A}, \cdot) .

Corollary

Let \mathfrak{g} be a solvable Lie algebra with an abelian complex structure J . Then:

- 1 \mathfrak{g}'_J is an abelian double product and if $\mathfrak{g}' \cap J\mathfrak{g}' = \{0\}$, then (\mathfrak{g}'_J, J) is holomorphically isomorphic to $\mathfrak{aff}(\mathcal{A})$ for some commutative associative algebra (\mathcal{A}, \cdot) ;
- 2 if $\mathfrak{g} = \mathfrak{g}' + J\mathfrak{g}'$, then $\mathfrak{g} = \mathfrak{u} \oplus J\mathfrak{u}$ is an abelian double product for some subalgebra $\mathfrak{u} \subset \mathfrak{g}'$.

Kähler Lie algebras with abelian complex structure

Let (\mathfrak{g}, J, g) be a Kähler Lie algebra with J abelian. It can be shown that:

- (i) $\mathfrak{z} = (\mathfrak{g}'_J)^\perp$.
- (ii) $(\mathfrak{g}')^\perp$ is abelian.
- (iii) $\text{ad}_z|_{\mathfrak{g}'}$ is symmetric for all $z \in \mathfrak{g}$.

Theorem (Andrada - B -Dotti, 2011)

Let (\mathfrak{g}, J, g) be a Kähler Lie algebra with J an abelian complex structure. Then \mathfrak{g} is isomorphic to

$$\mathfrak{aff}(\mathbb{R}) \times \cdots \times \mathfrak{aff}(\mathbb{R}) \times \mathbb{R}^{2s},$$

and this decomposition is orthogonal and J -stable.

Corollary

Let G be a simply connected Lie group equipped with a left-invariant Kähler structure (J, g) such that J is abelian. If the commutator subgroup is n -dimensional and the center is $2s$ -dimensional, then

$$G = H^2(-c_1) \times \cdots \times H^2(-c_n) \times \mathbb{R}^{2s},$$

where $c_i > 0$, $i = 1, \dots, n$, and $H^2(-c_i)$ denotes the 2-dimensional hyperbolic space of constant curvature $-c_i$.

Corollary

Let $M = \Gamma \backslash G$ be a compact quotient with a left invariant Kähler structure (J, g) such that J is abelian. Then M is diffeomorphic to a torus.

The first canonical Hermitian connection

Given a Hermitian Lie algebra (\mathfrak{g}, J, g) , consider the connection ∇^1 on \mathfrak{g} defined by

$$g(\nabla_x^1 y, z) = g(\nabla_x^g y, z) + \frac{1}{4}(d\omega(x, Jy, z) + d\omega(x, y, Jz)),$$

where ω is the Kähler form. This connection satisfies

$$\nabla^1 g = 0, \quad \nabla^1 J = 0, \quad T^1 \text{ is of type } (1, 1).$$

∇^1 is known as the first canonical Hermitian connection associated to (\mathfrak{g}, J, g) [Lichnerowicz, 1962].

Another expression for ∇^1 [Agricola, 2005]:

$$\nabla_x^1 y := \nabla_x^g y + \frac{1}{2} (\nabla_x^g J) Jy = \frac{1}{2} (\nabla_x^g y - J \nabla_x^g Jy),$$

for $x, y \in \mathfrak{g}$. We write the above equation with any affine connection ∇ and define

$$\bar{\nabla}_x y := \nabla_x y + \frac{1}{2} (\nabla_x J) Jy = \frac{1}{2} (\nabla_x y - J \nabla_x Jy), \quad (4)$$

for $x, y \in \mathfrak{g}$.

$\bar{\nabla}$ satisfies:

- $\bar{\nabla} J = 0$
- if ∇ is torsion-free, then $\bar{T}(x, y) = \bar{T}(Jx, Jy)$, i.e. \bar{T} is of type $(1, 1)$ with respect to J .

Lemma

Let ∇ be a torsion-free connection and J a complex structure on \mathfrak{g} . Assume that $\overline{\nabla} = 0$, that is, $\nabla_x J = -J\nabla_x$ for every $x \in \mathfrak{g}$. Then J is abelian.

Theorem (Andrada - B - Dotti, 2011)

Let (\mathfrak{g}, J, g) be a Hermitian Lie algebra such that its associated first canonical connection ∇^1 satisfies $\nabla_x^1 y = 0$ for every $x, y \in \mathfrak{g}$, that is, ∇^1 coincides with the $(-)$ -connection. Then \mathfrak{g} is abelian.

Corollary

Let $M = \Gamma \backslash G$ be a compact quotient of a simply connected Lie group G by a discrete subgroup Γ . If (J, g) is a left invariant Hermitian structure on M such that its first canonical connection ∇^1 coincides with the connection ∇^0 , then M is diffeomorphic to a torus.

Lemma

Let (\mathfrak{g}, J, g) be a Hermitian Lie algebra with J abelian. If the associated first canonical connection ∇^1 is flat, then $\mathfrak{z} \cap \mathfrak{g}' = \{0\}$.

Theorem

Let (\mathfrak{g}, J, g) be a Hermitian Lie algebra with J abelian. If the associated first canonical connection ∇^1 is flat, then \mathfrak{g} is abelian.