

BIFG

Boletín del Instituto de Fisiografía y Geología

ISSN 1666-115X

Volumen 92, 2023

Instituto de Fisiografía y Geología
Facultad de Ciencias Exactas, Ingeniería y Agrimensura
Universidad Nacional de Rosario

***Macroscaphites* (Ammonoidea)
in the Argille Scagliose Formation
of Prignano sulla Secchia
(Modena, Italy)**

Carlo Sarti



Boletín del Instituto de Fisiografía y Geología

ISSN 1666-115X

Fundado en 1937

Publicado por el Instituto de Fisiografía y Geología
"Dr. Alfredo Castellanos"
FCEIA - UNR - Argentina

Editorial

Instituto de Fisiografía y Geología
Facultad de Ciencias Exactas, Ingeniería, Agrimensura
Universidad Nacional de Rosario
Pellegrini 250, 2000 Rosario, Argentina

Correo-e: ifg@fceia.unr.edu.ar
Tel.: +00 54 341 4802649 (int. 135)
<http://www.fceia.unr.edu.ar/fisiografia/>

Editor

Horacio Parent - Universidad Nacional de Rosario,
Rosario

Comité editorial

Antonio Introcasa - Universidad Nacional de Rosario,
Rosario
Andrés F. Greco - Universidad Nacional de Rosario,
Rosario
Günter Schweigert - Staatliches Museum für
Naturkunde Stuttgart, Stuttgart
Alberto C. Garrido - Museo Olsacher, Dirección Provincial
de Minería Neuquén, Zapala
Gustavo González-Bonorino - Centro Austral de
Investigaciones Científicas, Ushuaia
Andrzej Kaim, Institute of Paleobiology - Polish Academy
of Sciences, Warsaw
Leonardo S. Filippi - Museo Municipal Argentino Urquiza,
Rincón de Los Sauces

Características y alcance

El Boletín IFG es una revista internacional de publicación periódica de artículos científicos originales sobre temas de las denominadas Ciencias de la Tierra (Geología, Estratigrafía, Paleontología, y disciplinas afines), cuyos tópicos estén o no incluidos en los campos de investigación del Instituto de Fisiografía y Geología. Sólo son aceptados manuscritos de trabajos inéditos consistentes en contribuciones científicas significativas a criterio de los Editores y los árbitros. Todos los manuscritos son sometidos a arbitraje internacional de por, al menos, dos reconocidos especialistas en la materia. La decisión final de aceptación de un manuscrito para su publicación es de los Editores.

Ediciones anteriores

El Boletín IFG es la serie única que reemplaza y sucede todas las ediciones anteriores del Instituto de Fisiografía y Geología que se publicaron con los nombres "Publicaciones" [ISSN 0041-8684], "Notas A" [ISSN 0325-4100] y "Notas B" [ISSN 0326-470X]. La numeración de la nueva y única serie comienza en el Volumen 70, correlativo con el último número de la antigua serie principal "Publicaciones", finalizada en 1996 con el número 69. Se publicará un volumen anual, dividido en un número variable de fascículos que dependerá de la extensión y número de contribuciones que se incluyan.

Suscripción e intercambio

El Boletín IFG puede obtenerse por intercambio institucional, suscripción individual o suscripción institucional, dirigiéndose a la dirección indicada arriba. La nómina de los números anteriores disponibles puede obtenerse del sitio de internet, así como todos los números anteriores disponibles en formato pdf.

Aim and scope

The Boletín IFG is an international periodical journal containing original scientific articles on the so called Earth Sciences (Geology, Stratigraphy, Paleontology, and related matters), related or not with the fields or research of the Instituto de Fisiografía y Geología. Only are accepted manuscripts of unpublished results consisting of significant scientific contributions. Every manuscript is subject to international peer review by at least two reviewers or referees, well known specialists on the matter. Final decision of publication is of the Editors.

Past editions

The Boletín IFG is the unique series of publications which replace and follows any former and past editions of the Instituto de Fisiografía y Geología which were published under the following names: "Publicaciones" [ISSN 0041-8684], "Notas A" [ISSN 0325-4100] y "Notas B" [ISSN 0326-470X]. The first volume of the series is the number 70 correlative with the last of the former series "Publicaciones", ended in 1996 with the number 69. It will be published one single volume per year, subdivided in a number of fascicles depending on the number and size of contributions included.

Suscription and exchange

The Boletín IFG can be obtained by institutional exchange, individual suscription or institutional suscription contacting to the adresses given above. All current and past publications are available on-line in the website cited above, in pdf format.

Macroscaphites (Ammonoidea) in the Argille Scagliose Formation of Prignano sulla Secchia (Modena, Italy)

Carlo Sarti



Recibido 15/03/2023
Aceptado 17/07/2023

Disponible vía Internet
10/08/2023
Versión impresa
10/12/2023

Editor: H. Parent

Sarti C., 2023. *Macroscaphites* (Ammonoidea) in the Argille Scagliose Formation of Prignano sulla Secchia (Modena, Italy). *Boletín del Instituto de Fisiografía y Geología* 92: 1-6. Rosario, 10/08/2023. ISSN 1666-115X.

Abstract. A new ammonite found in the “Argille Varicolori di Cassio” (Argille Scagliose Formation, tele-allochthonous Ligurides) of Prignano sulla Secchia, in Northern Apennines (Modena, Italy), is reported. The Argille Scagliose is a formation in which ammonites are practically absent. This is the first ammonite described found in the Modena region: an incomplete but exceptionally well-preserved specimen, which is taxonomically interpreted as a juvenile *Macroscaphites* cf. *tirolensis* Uhlig, 1888. The new finding allows dating the part of Argille Scagliose named “Argille Varicolori di Cassio” of the Modena region as Early Cretaceous (Early Barremian–earliest Late Barremian), whereas this unit was formerly considered Late Cretaceous. The hiatus reported in previous studies between Argille a Palombini Unit and the overlaying Argille Varicolori di Cassio Unit is therefore much shorter than previously thought.

Keywords: Heteromorph ammonite ▪ *Macroscaphites* cf. *tirolensis* ▪ Lower Barremian–basal Upper Barremian ▪ Argille Scagliose Formation ▪ Argille Varicolori di Cassio ▪ Prignano sulla Secchia.

Riassunto. Ritrovamento di una ammonite eteromorfa nelle Argille Scagliose di Prignano sulla Secchia, Modena:

Si segnala il ritrovamento di una ammonite appartenente alla specie *Macroscaphites* cf. *tirolensis* Uhlig, 1888, incompleta ma in eccezionale stato di conservazione, proveniente dalle “Argille Varicolori di Cassio” (Formazione delle Argille Scagliose, Liguridi tele-alloctone) di Prignano sulla Secchia (Modena). Nelle Argille Scagliose le ammoniti sono quasi del tutto assenti e in particolare questo è il primo ritrovamento descritto proveniente dal modenese. Le “Argille Varicolori di Cassio” erano finora considerate del Santoniano-Campaniano (Cretaceo superiore); il ritrovamento di *Macroscaphites* cf. *tirolensis* permette di datare questa porzione di successione, che si trova all'interno delle “Argille Scagliose”, al Barremiano inferiore-base del Barremiano superiore, quindi ci troviamo nella parte bassa del Cretaceo inferiore, e non nel Cretaceo superiore come erroneamente ritenuto fino ad oggi. La lacuna segnalata in letteratura tra le Argille a Palombini e la sovrastante Unità delle Argille Varicolori di Cassio (lacuna di quasi 60 milioni di anni) ora entrambe datate precisamente grazie al ritrovamento di due esemplari di ammoniti descritti da Sarti nel 2014 e nel presente lavoro è dunque molto più limitata di quanto supposto.

Parole chiave: Ammonite eteromorfa ▪ *Macroscaphites* cf. *tirolensis* ▪ Barremiano inferiore-base del Barremiano superiore ▪ Formazione delle Argille Scagliose ▪ Argille Varicolori di Cassio ▪ Prignano sulla Secchia.

Resumen. *Macroscaphites* (Ammonoidea) en la Formación Argille Scagliose de Prignano sulla Secchia (Módena, Italia): En el presente trabajo se describe un nuevo amonite descubierto en la unidad “Argille Varicolori di Cassio” (Formación Argille Scagliose, Ligurides tele-allochthonas) de Prignano sulla Secchia, en los Apenninos del Norte (Módena, Italia). En la Formación Argille Scagliose los amonites son prácticamente ausentes. El amonite descrito en este trabajo es el primero hallado en la región de Módena: un ejemplar incompleto pero excepcionalmente preservado que se interpreta taxonómicamente como un *Macroscaphites* cf. *tirolensis* Uhlig, 1888 juvenil. Este nuevo hallazgo permite la datación de la Argille Scagliose named “Argille Varicolori di Cassio” en la región de Modena como Cretácico Temprano (Barremiano Temprano-Late Barremian), en tanto esta unidad era considerada Cretácico Tardío. El hiato establecido en estudios anteriores entre la unidad Argille a Palombini y la sobreyacente unidad Argille Varicolori di Cassio es por lo tanto mucho mas breve.

Palabras clave: Amonite heteromorfo ▪ *Macroscaphites* cf. *tirolensis* ▪ Barremiano inferior-Barremiano superior basal ▪ Formación Argille Scagliose ▪ Argille Varicolori di Cassio ▪ Prignano sulla Secchia.

Carlo Sarti [carlo.sarti@unibo.it]: Collezione Museo di Geologia e Paleontologia “Giovanni Capellini”, Università di Bologna, Via Zamboni 63-67, 40127 Bologna, Italy.

AMMONITE RECORDS FROM THE ARGILLE SCAGLIOSE FORMATION (NORTHERN APENNINES)

The ammonites found in the Argille Scagliose Formation (Bianconi 1840) are very rare but very important for precise dating of this unit. The first report of ammonites from the Argille Scagliose Formation was given by the famous naturalist Antonio Stoppani (1875): a specimen from Costa de Grassi, near Reggio Emilia. Two years later, two ammonite specimens were collected by Mantovani (1877) in the Argille Scagliose of Ranzano (Parma) and Monte dell'Evangelo near Scandiano (Reggio Emilia).

In his paper: "Contribution à la Connaissance paléontologique des Argiles écaillées" Federico Sacco (1893) described a fauna with ammonites, and concluded that ammonites in the Argille Scagliose are Cretaceous. Sacco (1893) described ammonites of the genera *Hamites* (4 fragmentary and poorly preserved specimens from Valle Del Dordone, one of them probably an ichnofossil), *Acanthoceras* (2 poorly preserved specimens from Cà di Panico in Val Marecchia), *Hoplites* (3 fragmentary specimens from Ranzano in Val d'Enza and Val di Taro), *Desmoceras* (2 well preserved specimens from Valle del Dordone), and *Pachydiscus* (2 ammonites).

In a second paper Sacco (1905) described ammonites assigned to the genera *Hamites*, *Scaphites*, *Toxoceras*, *Ancyloceras*, *Baculites*, *Turrilites*, *Desmoceras*, and *Acanthoceras*.

Finally, Sacco (1923) returned to the subject and reported *Hamites* from Pellegrino Parmense (Parma), and a specimen from Val Dordone (near Parma) assigned to the genus *Schloenbachia*.

Anelli (1935, 1938) collected some ammonites in the Argille Scagliose of the Emilia Apennines. Due to the poor preservation of the material, identifications were difficult. Notable is only a well preserved specimen of *Crioceras*, from Cervarezza (Reggio Emilia Apennines).

To my knowledge, later findings of ammonites were no longer reported from the Argille Scagliose. However, all these historical findings do not provide detailed stratigraphic information. This is because of the inaccurate information about the localities of provenance, and because lithological descriptions were insufficient, even for approaching a broad-stratigraphic correlation (e.g., *fide* Sacco 1905: *Acanthoceras* is Middle Cenomanian, as *Turrilites*, but *Baculites* is Upper Turonian or younger). Despite the fact that the locality data are not well-known an update of the identifications in a paper in preparation will be of much help.

Recently a very interesting specimen of *Pulchellia* (*Pulchellia*) n. sp. aff. *kiliani* Hyatt, 1903 has been described by Sarti (2014). This specimen comes from the Barremian of Argille Scagliose exposed near the East flank of Mount Montovolo (Province of Bologna), in front of the stratigraphic section Montovolo Sud, in the part of Argille Scagliose named Palombini shales (tele-allochthonous Ligurides, see Vai & Martini 2001). This specimen is the



Figure 1. A: Location of the study area. B: Topographic map of the study area with indication of the sampling point (red dot).

first ammonite described from the part of Argille Scagliose, named Palombini shales. It is indeed a very important finding because of its very good preservation and its accurate stratigraphic location. The Palombini Shales represent one of the most typical and widespread units cropping-out in the Northern Apennines, and in the Ligurid sequences (oceanic sequences) these are part of the ophiolite sequence sedimentary cover (Cobianchi & Villa 1992).

The finding of the ammonite *Macroscaphites* cf. *tirolensis* Uhlig, 1888 described in the present paper, is very important because of its accurate stratigraphic position. This ammonite indicates a stratigraphic position in the interval Lower Barremian to base of the Upper Barremian, making possible for the first time to date by an ammonite the part of the Argille Scagliose of the Emilian Apennines named “Argille Varicolori di Cassio” (Fig. 1).

STRATIGRAPHIC FRAMEWORK

The ammonite described in this paper was discovered by the collector Giovanni Andreoli in the Argille Scagliose Fm of Prignano sulla Secchia (Modena), in the Caselletta Quarry, where it is mainly exploited by reddish clays materials for ceramic and brick industry in the “Argille rosse della Val Rossenna”. The clays are mined from blocks embedded in the Argille Varicolori (MVRb of the Geological Map of Italy, Gasperi et al. 2005). There are also clays that are being mined in the “Argille Varicolori di Cassio” Unit (AVV) that certainly look very similar to the “Argille rosse della Val Rossenna”, and they could be confused. However, their ages are very different, the former is Cenozoic, while the Argille Varicolori di Cassio is Mesozoic. In fact, regarding the Argille rosse della Val Rossenna Unit, it is worth to note that the age is Eocene, perhaps Maastrichtian the base of the Unit MVRb (Gasperi et al. 2005). The present ammonite discovered in the Caselletta Quarry suggests that the “Argille rosse” could be attributed at least partially to the “Argille Varicolori di Cassio”, that is a Cretaceous Unit whose lower boundary is the Argille a Palombini.



Figure 2. *Macroscaphites* cf. *tirolensis* Uhlig, 1888, Argille Varicolori di Cassio of the Argille Scagliose Formation, teleallochthonous Ligurides. Caselletta Quarry, Prignano sulla Secchia (Modena, Italy), Lower Barremian-basal Upper Barremian. – Double size. The scale bar represents 10 mm.

The thickness of the Argille Varicolori di Cassio is about 200 meters. On the basis of published data (Rio & Villa 1987, Gasperi et al. 2005, Serpagli 2005, Barbero et al. 2017, Serafini et al. 2019, among others) the unit ranges from the Santonian up to the Campanian. This is suggested by the analyses of microfossil associations that are attributed to the Campanian with doubts, and from ichnofossils and macrofossils (mostly Gastropoda, Lamellibranchia and Ichthyosaurs) which are poorly significant for stratigraphic purposes. Considering the present specimen of *Macroscaphites* cf. *tirolensis*, the age of the Argille Varicolori di Cassio is now fixed as Early Cretaceous, not Late Cretaceous as previously reported by former authors. The known chronostratigraphic range of *Macroscaphites tirolensis* is the interval Early Barremian-basal Late Barremian.

The hiatus reported in previous studies between the Argille a Palombini Unit and the overlying Argille Varicolori di Cassio Unit (about 60 million of years!), is much reduced after the ammonite record of Sarti (2014) and that of the present paper.

Table 1. Measurements (in mm) of the studied and comparative specimens. Abbreviations: *D*: shell diameter, *H*: whorl height, *U*: umbilical diameter, *W*: whorl width, *N.ribs*: number of primary ribs per whorl, *N/2*: number of primary ribs per half whorl.

	<i>D</i>	<i>H</i>	<i>U</i>	<i>W</i>	<i>H/D</i>	<i>U/D</i>	<i>W/D</i>	<i>N.ribs</i>
<i>Macroscaphites</i> cf. <i>tirolensis</i> (Fig. 2)	26.4	8.4	12.0	~7.0	0.32	0.45	~0.26	99-100
	20.4	6.3	8.8	-	0.31	0.40	-	84-85
	14.0	5.3	6.0	~5.5	0.38	0.43	~0.39	~ 70
	12.0	3.6	4.8	-	0.30	0.40	-	62-63
	10.5	3.1	4.5	-	0.30	0.43	-	52-53
<i>Macroscaphites tirolensis</i> (Holotype)	33.0	10.0	17.0	~6.0	0.30	0.51	0.18	99-100
	26.0	7.0	13.0	-	0.27	0.50	-	80-81
<i>Macroscaphites yvani</i> (Holotype)	55.5	13.8	30.1	12.0	0.25	0.54	0.22	84
	41.9	11.8	22.6	-	0.28	0.54	-	82
	~ 26	~ 7.5	~ 13	-	~0.29	~0.5	-	~ 68
<i>Macroscaphites yvani</i> (Baudoin et al. 2012: 632; crn60)	49.3	14.1	24.0	-	0.29	0.49	-	47 (N/2)
	42.3	13.4	19.7	-	0.32	0.47	0.37	49 (N/2)
	37.9	12.6	17.2	-	0.33	0.45	0.38	50 (N/2)
	31.3	10.7	14.2	-	0.34	0.45	0.41	-
	25.4	7.3	12.3	-	0.29	0.48	0.45	-
	19.4	5.2	8.5	-	0.27	0.44	-	-




Figure 3. A: Sketch (modified) from Avram (1984: fig. 8b) of the holotype of *Macroscaphites tirolensis*. B: The specimen of *Macroscaphites* cf. *tirolensis* described in the present paper (slightly reduced), superimposed on the sketch A for comparison.

SYSTEMATIC PALAEONTOLOGY

The specimen described was donated by Giovanni Andreoli to the Museum of Ecology and Natural History of Marano sul Panaro (Modena): inventory number 901/GF.

Superfamily Ancyloceratoidea Gill, 1871

Family Macroscaphitidae Hyatt, 1900

Genus *Macroscaphites* Meek, 1876

Type species. *Scaphites yvani* Puzos, 1832; subsequent designation by Roman (1938: 38).

Remarks. Since Munier-Chalmas (1892) *Macroscaphites* and *Costidiscus* Uhlig, 1882 are considered possible dimorphic partners (Avram 1984). Among the Barremian species of *Macroscaphites* and *Costidiscus* a few constitute surely dimorphic pairs, controlled by the accurate recording, bed by bed, of the material, e.g. *Macroscaphites yvani* and *Costidiscus recticostatus* (d'Orbigny, 1841), see Baudouin et al. (2012). Some other species were grouped in

dimorphic pairs with doubts and others have not presumable partners currently. This is the case of the microconchiate *M. tirolensis*.

***Macroscaphites* cf. *tirolensis* Uhlig, 1888**

Figs. 2-4

cf. 1888 *Macroscaphites tirolensis* n. sp. – Uhlig: 86, pl. 4: 2.

cf. 1920 *Macroscaphites tirolensis* Uhlig. – Gignoux: 117.

cf. 1972 *Macroscaphites?* *tirolensis* Uhlig. – Vasicek: 48, pl. 3: 2.

cf. 1984 *Macroscaphites tirolensis* Uhlig. – Avram: 73, fig. 8b.

Material. One specimen with inventory number 901/GF.

Locality of provenance. Caselletta Quarry, Prignano sulla Secchia (Modena, Italy).

Stratigraphic location. “Argille Varicolori di Cassio” (Argille Scagliose Fm, tele-allochthonous Ligurides).

Description. Small, very evolute microconch (maximum diameter about 26.5 mm), planispirally coiled with slightly covering whorls, with broad and shallow umbilicus and gentle, low umbilical wall. Whorl section slightly depressed-oval, whorl width-to-high ratio (W/H) = 1.03 at 14 mm diameter (D), becoming subcircular (W/H = 0.95 at D = 26 mm) with venter slightly convex. Ornamentation consists of dense straight and radial ribs in the inner whorls, becoming slightly prorsiradiate on the outer half-whorl. Ribs mainly simple, very rarely bifurcated at mid-flank. The ribbing is not uniform, increasing in density through the ontogeny: 52-53 ribs per whorl at D = 10.5 mm diameter, 70 at D = 14 mm, to 99-100 at D = 26.4 mm. The ribs are well marked and sharp on the inner whorls, but weaker and rounded in the last whorl, and crossing the venter apparently uninterrupted. On the last preserved part the shell shows two rows of nodes, one close to the umbilicus and the second at the transition of the flanks to the venter. At the diameters of approximately 26 mm, 25 mm and 24.6 mm, three ribs show tubercles elongated, small at the transition of umbilicus to flanks, and more prominent at the transition of the flanks to the venter. These are well marked, forming ventro-lateral claviform nodes slightly prorsiradiate, and after they broaden out, forming three slightly thicker and more evident ribs on the venter. One rib, slightly more pronounced, with tubercles occurs between 3 to 4 simple ribs. Only one constriction occurs in the inner whorls (D = 9 mm). The constriction is narrow and parallel to the ribs. The specimen preserves only the spiral whorls of the phragmocone. The typical straight or curved shaft and recurved crozier body chamber is not preserved (Fig. 3).

Discussion. It is not possible to make an exact specific assignment of the ammonite because the final part of the coiled and the entire uncoiled part are not preserved. The specimen matches the morphological characteristics of the species *Macroscaphites tirolensis* Uhlig, 1888 but with slight differences with the holotype: narrower umbilicus, greater whorl width, and partly whorl height, and, moreover shows a lower numbers of ribs.

It differs from *Macroscaphites binodosus* Uhlig, 1883 by the slightly smaller size and more prominent ribs. Tubercles or nodes are elongated in the present species but rounded in *M. binodosus*.

Macroscaphites olcostephanooides (Uhlig, 1883) differs from *M. tirolensis* by its ribs forming bullae only on the umbilical border, and bifurcating in the inner whorls, as well as by its more involute coiling.

M. yvani differs from *M. tirolensis* by the complete lack of tubercles through the ontogeny; our specimen bears small umbilical nodes and well marked ventro-lateral claviform tubercles on the last preserved part of the shell. Furthermore, the relative umbilical width is larger in the holotype of *M. yvani*. Another difference of our specimen with respect to *M. yvani* is the lower number of ribs at comparable diameter (Fig. 4). A species similar to *M. yvani* is *Macroscaphites striatisulcatus* (d'Orbigny, 1841) which differs from *M. tirolensis* by its much more frequent bifurcate ribs and more frequent constrictions.

Stratigraphic distribution. *M. tirolensis* is recorded only in the Barremian (see Vasicek 1972) and has been mentioned with accurate stratigraphical position from the




Figure 4. *Macroscaphites* cf. *tirolensis* Uhlig, 1888. Primary rib-density curves of the specimen described in this paper and species discussed in text. (1) *Macroscaphites* cf. *tirolensis* Uhlig, 1888; (2) *Macroscaphites yvani* (Puzos, 1832) in Baudouin et al. (2012: pl. 9: 1); (3) *Macroscaphites yvani* (Puzos, 1832) in Baudouin et al. (2012: pl. 9: 3); (4) *Macroscaphites yvani* (Puzos, 1832) in Baudouin et al. (2012: pl. 9: 2); (5) *Macroscaphites yvani* (Puzos, 1832), holotype; (6) *Macroscaphites tirolensis* Uhlig, 1888, holotype; (7) *Macroscaphites binodosus* Uhlig, 1883, holotype.

lower Barremian, base of the Moutonianum Zone (corresponding to the base of the Darsi Zone) in the Puez area, South Tyrol, North Italy (Lukeneder & Lukeneder 2014), middle-upper parts of the Darsi Zone (Vermeulen 1997, 1999) and base of the upper Barremian in the Carpathian area (Avram 1983).

CONCLUSIONS

Known fossils from the Argille Scagliose Formation are very rare, including ammonites. The few references found in literature are almost all from the mid 19th to the early 20th century. The recently described *Pulchellia* n. sp. aff. *kiliani* is remarkable by the accurate stratigraphic location of the specimen (Sarti 2014). The relevance of the finding of the exceptionally well preserved specimen of *Macroscaphites* cf. *tirolensis* described in the present paper, is remarkable by its stratigraphic location, allowing a new understanding on the chronostratigraphic position as Lower Barremian-basal Upper Barremian that, for the first time with an ammonite, makes it possible to date the part of the Argille Scagliose of the Emilian Apennines named “Argille Varicolori di Cassio”.

Acknowledgements. I would like to thank Dr. Prof. Jens Lehmann (University of Bremen, Germany) for his useful suggestions and assistance in reviewing this paper. I am also indebted to Dr. Prof. Gerard Delanoy (University of Nice, France) for the careful review of this paper and helpful comments. I am grateful to Giovanni Andreoli for granting access to the ammonite and in providing the topographic map of the study area.

REFERENCES

- Anelli M., 1935. Appunti paleontologici a proposito delle cosiddette "Argille Scagliose". – *Rivista Italiana di Paleontologia* **41**: 33-44.
- Anelli M., 1938. Sulla presenza di Aptici nelle cosiddette argille scagliose dell'Appennino emiliano. – *Rivista Italiana di Paleontologia* **44**: 82-93.
- Avram E., 1983. Barremian ammonite zonation in the Carpathian area. – *Zitteliana* **10**: 509-514.
- Avram E., 1984. Correspondent species of the genera *Macroscaphites* Meek and *Costidiscus* Uhlig. – *University of Bucharest, Laboratory of Paleontology Special Volume*: 67-80.
- Baudouin C., Delanoy G., Boselli P., Bert D. & Boselli M., 2012. Les faunes d'ammonites de la sous-zone à Sarasini (Barrémien supérieur) dans les Baronnies (Drôme, France). – *Revue de Paléobiologie* **31**: 601-677.
- Barbero E., Festa A., Fioraso G. & Catanzariti R., 2017. Geology of the Curone and Staffora Valleys (NW Italy): field constraints for the Late Cretaceous-Pliocene tectono-stratigraphic evolution of Northern Apennines. – *Journal of Maps* **13**: 879-891.
- Bianconi G.G., 1840. Storia naturale dei terreni ardenti, dei vulcani fangosi, delle sorgenti infiammabili dei pozzi idropirici e di altri fenomeni geologici operati dal gas idrogeno e della origine di esso gas, del Dott. Giuseppe Bianconi. – Jacopo Marsigli Editore, Bologna, 216 pp.
- Cobianchi M. & Villa G., 1992. Biostratigrafia del calcare a Calpionelle e delle Argille a Palombini nella sezione di Statale (Val Graveglia, Appennino ligure). – *Atti Ticinesi Scienze della Terra* **35**: 199-211.
- Gasperi G., Bettelli G., Panini F. & Pizzio M., 2005. *Note illustrative della carta geologica d'Italia alla scala 1:50.000, foglio 219 Sassuolo*: 1-195.
- Gignoux M., 1920. Les Phyllocératidés, Lytocératidés et Pulchelliidés du Paléocrétacé. In: W. Kilian (ed.): Contribution à l'étude des Céphalopodes paléocrétacés du SE de la France. – *Mémoires pour servir à l'explication de la carte géologique détaillée de la France*: 85-165.
- Hyatt A., 1900. Cephalopoda. In: Eastman's Zittel textbook of Paleontology, First edition, pp. 502-604. New York.
- Hyatt A., 1903. Pseudoceratites of the Cretaceous. *United States Geological Survey Monograph* **44**: 128-144.
- Kakabadze M.V. & Thieuloy J.P., 1991. Ammonites hétéromorphes du Barrémien et de l'Aptien de Colombie, Amérique du Sud. – *Géologie Alpine* **67**: 81-113.
- Lukeneder A. & Lukeneder S., 2014. The Barremian heteromorph ammonite *Dissimilites* from northern Italy: Taxonomy and evolutionary implications. – *Acta Palaeontologica Polonica* **59**: 663-680.
- Mantovani P., 1877. Intorno ad alcuni ammonite dell'Appennino dell'Emilia. – Tipografia degli Artigianelli, Reggio Emilia: 1-13.
- Meek F.B., 1876. A report on the invertebrate Cretaceous and Tertiary fossils of the upper Missouri country. – *Report of the United States Geological Survey of the Territories* **9**: 1-629.
- Orbigny d'A., 1840-1842. Paléontologie française, description zoologique et géologique de tous les animaux mollusques et rayonnés fossiles de France. Terrains Crétacés, I. Céphalopodes. – Masson éd., Paris, 662pp.
- Puzos M., 1832. Sur le *Scaphites* Yvoni. – *Bulletin de la Société Géologique de la France* **2**: 355-356.
- Rio D. & Villa G., 1987. On the age of the "Salti del Diavolo" conglomerates and of the Monte Cassio Flysch "Basal Complex" (Northern Apennines, Parma Province). – *Giornale di Geologia* **3**: 63-79.
- Roman F., 1938. Les ammonites jurassiques et crétacées. Essai de genera. – Masson, Paris, 554 pp.
- Sacco F., 1893. Contribution à la connaissance paléontologique des argiles écaillées et des Schistes ophiolitiques de l'Apennin septentrional. – *Bulletin de la Société belge de géologie, de paléontologie et d'hydrologie* **7**: 3-34.
- Sacco F., 1905. Les Formations ophitiformes du Crétacé. – *Bulletin de la Société belge de géologie, de paléontologie et d'hydrologie* **19**: 247-265.
- Sacco F., 1923. Nuovi fossili Cretacei negli argilloschisti dell'Appennino settentrionale. – *Atti Reale Accademia delle Scienze di Torino* **54**, **247**: 30-43.
- Sarti C., 2014. New find of an Ammonite in the "Argille Scagliose" Formation of Montovolo (Bologna). – *Neues Jahrbuch für Geologie und Paläontologie* **274**: 127-132.
- Serafini G., Rondelli R., Fornaciari B. & Papazzoni C.A., 2019. Segnalazione di un nuovo ittiosauro dal Cretaceo dell'Appennino modenese (Pavullo nel Frignano). – *Atti della Società dei naturalisti e matematici di Modena* **150**: 111-120.
- Serpagli E., 2005. First record of the ichnofossil *Atollites* from the Late Cretaceous of the Northern Apennines, Italy. – *Acta Palaeontologica Polonica* **50**: 403-408.
- Stoppani A., 1875. Delle Argille scagliose e di alcuni Ammoniti dell'Appennino dell'Emilia. – *Atti della Società Italiana di Scienze Naturali* **18**: 1-19.
- Uhlig V., 1883. Die Cephalopodenfauna der Wernsdorfer Schichten. – *Denkschriften der Kaiserlichen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche* **46**: 127-290.
- Uhlig V., 1888. Ueber neocom Fossilien von Gardenazza in Südtirol, nebst einem Anhang über das Neocom von Ischl. – *Jahrbuch der Kaiserlich Königlich Geologischen Reichsanstalt* **37**: 69-108.
- Vai G.B. & Martini I.P., 2001. Anatomy of an Orogen. The Apennines and adjacent Mediterranean Basins. – Kluwer Academic Publisher, The Netherlands, 639 pp.
- Vašíček Z., 1972. Ammonoidea of the Těšín-Hradiště Formation (Lower Cretaceous) in the Moravskoslezské Beskydy Mts. – *Rozprawy Ústředního Ústavu Geologického* **38**: 3-137.
- Vermeulen J., 1997. Biohorizons ammonitiques dans le Barrémien du Sud-Est de la France (de la zone à Hugii à la zone à Sartousiana). – *Géologie Alpine* **73**: 99-117.
- Vermeulen J., 1999. Nouvelles données sur les répartitions stratigraphiques, les évolutions et les classifications de trois familles d'ammonites du Crétacé inférieur. – *Géologie Alpine* **75**: 123-132.
- Vermeulen J., 2005. Boundaries, ammonite fauna and main subdivisions of the stratotype of the Barremian. In: T. Adatte, A. Thierry, A. Godet, A. Arnaud-Vanneau, H. Arnaud & K. Foellmi (eds.): The Hauterivian-Lower Aptian Sequence Stratigraphy from Jura Platform to Vocontian Basin: A Multidisciplinary Approach. – *Géologie Alpine, série spéciale «colloques et excursions»* **7**: 147-173.

BIFG

Boletín del Instituto de Fisiografía y Geología
ISSN 1666-115X

Volumen 92, 2023

Rosario - 2023